

# ON A CLASS OF $H$ -SELFADJOINT RANDOM MATRICES WITH ONE EIGENVALUE OF NONPOSITIVE TYPE.

MICHAŁ WOJTYŁAK

**ABSTRACT.** Large  $H$ -selfadjoint random matrices are considered. The matrix  $H$  is assumed to have one negative eigenvalue, hence the matrix in question has precisely one eigenvalue of nonpositive type. It is showed that this eigenvalue converges in probability to a deterministic limit. The weak limit of distribution of the real eigenvalues is investigated as well.

## INTRODUCTION

The main object of this survey are random matrices that are not symmetric, but are selfadjoint with respect to an indefinite inner product. Spectrum and numerical range of some classes of such matrices were considered recently in [23, 24], although, in the present paper we consider different instances. The infinite linear algebra motivation of the present research is the following. Consider an invertible, hermitian-symmetric matrix  $H \in \mathbb{C}^{n \times n}$ . We say that  $X \in \mathbb{C}^{n \times n}$  is  $H$ -selfadjoint if  $X^*H = HX$ . This is the same as to say that  $A$  is selfadjoint with respect to an inner product

$$[x, y]_H := y^* H x, \quad x, y \in \mathbb{C}^n.$$

Note that this inner product is not positive definite if  $H$  has negative eigenvalues. In the literature the space  $\mathbb{C}^n$  with the inner product  $[\cdot, \cdot]_H$  is also called  $\Pi_\kappa$ -space (where  $\kappa$  is the number of negative eigenvalues of  $H$ ) or Pontryagin space, the infinite dimensional case is considered as well [5, 20]. In the present paper the case when

$$(1) \quad H = \begin{bmatrix} -1 & 0 \\ 0 & I_N \end{bmatrix},$$

is considered. It is easy to check that for such  $H$  each  $H$ -selfadjoint matrix has the form

$$(2) \quad X = \begin{bmatrix} a & -b^* \\ b & C \end{bmatrix},$$

with  $x \in \mathbb{R}$ ,  $b \in \mathbb{C}^N$  and a hermitian-symmetric matrix  $C \in \mathbb{C}^{N \times N}$ . Due to the famous theorem of Pontryagin [27] the matrix  $X$  has precisely one eigenvalue  $\beta$ , for which the corresponding eigenvector  $x$  satisfies  $[x, x]_H \leq 0$ . The problem of tracking the nonpositive eigenvalue was considered for example in [9, 29]. In those papers the setting was non-random and  $X$  was in the family of one dimensional

---

2000 *Mathematics Subject Classification.* Primary 15B52, Secondary 15B30, 47B50.

*Key words and phrases.* Random matrix, Wigner matrix, eigenvalue, limit distribution of eigenvalues,  $\Pi_1$ -space.

The research was supported by Polish Ministry of Science and Higher Education with a Iuventus Plus grant.




FIGURE 1. Eigenvalues of a random matrix  $X_{100}$  computed with R [28]

extensions of a fixed operator in an infinite dimensional  $\Pi_1$ -space. The aim of the present work is to investigate the behavior of  $\beta$  when  $X$  is a large random matrix. We show that the main method of [9, 29] – the use of Nevanlinna functions with one negative square – can be adapted to the random setting as well.

A classical result of Wigner [31] says that if the random variables  $y_{ij}$ ,  $0 \leq i \leq j < +\infty$  are real, i.i.d with mean zero and variance equal one, then the distribution of eigenvalues of a matrix

$$Y_N = \frac{1}{\sqrt{N}} [y_{ij}]_{i,j=0}^N,$$

where  $y_{ji} = y_{ij}$  for  $j > i$ , converge weakly in probability to the Wigner semi-circle measure. Note that by multiplying the first row of  $Y_N$  by -1 we obtain a  $H$ -selfadjoint matrix  $X_N$ . A result of a preliminary numerical experiment with gaussian  $y_{ij}$  is plotted in Figure 1. Note that the spectrum of  $X_N$  is real, except two eigenvalues, lying symmetrically with respect to the real line. Although we pay a special attention to the above case, we study the behavior of the eigenvalue of nonpositive type in a more general setting. Namely, we assume that the random matrix  $X_N$  in  $\mathbb{C}^{N \times N}$  is of a form

$$X = \begin{bmatrix} a_N & -b_N^* \\ b_N & C_N \end{bmatrix},$$

with  $a_N$ ,  $b_N$  and  $C_N$  being independent. Furthermore, the vector  $b_N$  is a column of a Wigner matrix and  $a_N$  converges weakly to zero. The only assumption on  $C_N$  is that the limit distribution of its eigenvalues converge weakly in probability, see (R0)–(R3) for details. In Theorem 6 we prove that under these assumptions the non-real eigenvalues converge in probability to a deterministic limit that can be computed knowing the limit distribution of eigenvalues of  $C_N$ . In the case when  $C_N$  is a Wigner matrix the nonreal eigenvalues converge to  $\pm i\sqrt{2}/2$ , cf. Theorem 8. Furthermore, under a technical assumption of continuity of the entries of  $X_N$ , we show in Theorem 7 that the limit distribution of the real eigenvalues of  $X_N$

coincides with the limit distribution of eigenvalues for the matrices  $C_N$ . Again, in the case when  $C_N$  is a Wiegner matrix we obtain a more precise result. Namely, in Theorem 8 we show that the real eigenvalues  $\zeta_2^N, \dots, \zeta_N^N$  of  $X_N$  and the eigenvalues  $\lambda_1^N, \dots, \lambda_N^N$  of  $C_N$  satisfy the following inequalities:

$$\lambda_1^N < \zeta_2^N < \lambda_2^N < \dots < \lambda_{N-1}^N < \zeta_N^N < \lambda_N^N.$$

It shows that the nonreal eigenvalue of  $X_N$  plays an analogue role as the largest eigenvalue in one-dimensional, symmetric perturbations of Wiegner matrices. This fact relates the present paper to the current work on finite dimensional perturbations of random matrices, see [2, 3, 4, 6, 7, 13, 15] and references therein. Also note that  $X_N$  is a product of a random and deterministic matrix, such products were already considered in the literature, see e.g. [30].

The author is indebted to Anna Szczepanek for preliminary numerical simulations and Maxim Derevyagin for his valuable comments. Special thanks to Anna Kula, Patryk Pagacz and Janusz Wysoczyński.

## 1. FUNCTIONS OF CLASS $\mathcal{N}_1$

The Nevanlinna functions with negative squares play a similar role for the class of  $H$ -selfadjoint matrices as the class of ordinary Nevanlinna function plays for hermitian-symmetric matrices. This phenomenon has its roots in operator theory, we refer the reader to [8, 9, 16, 17, 18] and papers quoted therein for a precise description of a relation between  $\mathcal{N}_\kappa$ -functions and selfadjoint operators in Krein and Pontryagin spaces. We begin with a very general definition of the class  $\mathcal{N}_\kappa$ , but we immediately restrict ourselves to certain subclasses of those functions.

We say that  $Q$  is a *generalized Nevanlinna function of class  $\mathcal{N}_\kappa$*  [16, 19] if it is meromorphic in the upper half-plane  $\mathbb{C}^+$  and the kernel

$$N(z, w) = \frac{Q(z) - \overline{Q(w)}}{z - \bar{w}}$$

has precisely  $\kappa$  negative squares, that is for any finite sequence  $z_1, \dots, z_k \in \mathbb{C}^+$  the hermitian-symmetric matrix

$$[N(z_i, z_j)]_{i,j=1}^k$$

has not more than  $\kappa$  nonpositive eigenvalues and for some choice of  $z_1, \dots, z_k$  it has precisely  $\kappa$  nonpositive eigenvalues. In the present paper we use this definition with  $\kappa = 0, 1$ .

The class  $\mathcal{N}_0$  is the class of ordinary Nevanlinna functions, i.e. the functions that are holomorphic in  $\mathbb{C}^+$  with nonnegative imaginary part. By  $M_b^+(\mathbb{R})$  we denote the set of positive, bounded Borel measures on  $\mathbb{R}$ . For  $\mu \in M_b^+(\mathbb{R})$  we define the Stieltjes transform as

$$\hat{\mu}(z) = \int_{\mathbb{R}} \frac{1}{t - z} dt, \quad z \in \mathbb{C} \setminus \text{supp } \mu.$$

Clearly,  $\hat{\mu}$  belongs to the class  $\mathcal{N}_0$  and the values of  $\hat{\mu}$  in the upper half-plane determine the measure uniquely by the Stieltjes inversion formula. Although not every function of class  $\mathcal{N}_0$  is a Stieltjes transform of a Borel measure (cf. [11]), this subclass of  $\mathcal{N}_0$  functions will be sufficient for present reasonings. Also, we will be interested in a special subclass of  $\mathcal{N}_1$  functions, namely in the functions of the form (3) below. We refer the reader to the literature [8, 10] for representation theorems for  $\mathcal{N}_\kappa$  functions.

**Proposition 1.** *If  $\mu \in M_b^+(\mathbb{R})$ ,  $a \in \mathbb{R}$  then*

$$(3) \quad Q(z) = \hat{\mu}(z) + a - z$$

*is a holomorphic function in  $\mathbb{C}^+$  and belongs to the class  $\mathcal{N}_1$ . Furthermore, there exists precisely one  $z_0 \in \mathbb{C}$  such that either  $z_0 \in \mathbb{C}^+$  and*

$$(4) \quad Q(z_0) = 0,$$

*or  $z_0 \in \mathbb{R}$  and*

$$(5) \quad \lim_{z \rightarrow z_0} \frac{Q(z)}{z - z_0} \in (-\infty, 0].$$

The symbol  $\rightarrow$  above denotes the non-tangential limit:

$$z \in \mathbb{C}^+, \quad z \rightarrow z_0, \quad \pi/2 - \theta \leq \arg(z - z_0) \leq \pi/2 + \theta,$$

with some  $\theta \in (0, \pi/2)$ . We call  $z_0 \in \mathbb{C}^+ \cup \mathbb{R}$  the *generalized zero of nonpositive type (GZNT)* of  $Q(z)$ . The first part of the Proposition can be found e.g. in [17], while for the proof of the 'Furthermore' part in the general context<sup>1</sup> we refer the reader to [19, Theorem 3.1, Theorem 3.1']. In view of the above proposition we can define a function

$$G : M_b^+(\mathbb{R}) \times \mathbb{R} \rightarrow \mathbb{C}^+$$

by saying that  $G(\mu, a)$  is the GZNT of the function  $\hat{\mu}(z) + a - z$ . The following proposition plays a crucial role in our arguments.

**Proposition 2.** *The function  $G$  is jointly continuous with respect to the weak topology on  $M_b^+(\mathbb{R})$  and the standard topology on  $\mathbb{R}$ .*

*Proof.* Assume that  $(\mu_n)_n \subset M_b^+(\mathbb{R})$  converges weakly to  $\mu \in M_b^+(\mathbb{R})$  and  $a_n \in \mathbb{R}$  converges to  $a \in \mathbb{R}$  with  $n \rightarrow \infty$ . Take a compact  $K$  in the open upper half-plane, with nonempty interior. Then  $\hat{\mu}_n$  converges uniformly to  $\hat{\mu}$  on the set  $K$ . Indeed, if  $r = \sup_{t \in \mathbb{R}, z \in K} 1/|t - z|$  then

$$\sup_{z \in K} |\hat{\mu}_n(z) - \hat{\mu}_0(z)| \leq r |\mu_n - \mu_0|(\mathbb{R}),$$

the latter clearly converging to zero with  $n \rightarrow \infty$ . In consequence,  $\hat{\mu}_n(z) + a_n - z$  converges to  $\hat{\mu}(z) + a - z$  uniformly on  $K$  with  $n \rightarrow \infty$ . By [21] the GZNT of  $\hat{\mu}_n(z) + a_n - z$  converges to the GZNT of  $\hat{\mu}(z) + a - z$ , which finishes the proof.  $\square$

## 2. $H$ -SELFADJOINT MATRICES

In this section we review basic properties of selfadjoint matrices in indefinite inner product spaces introducing the concept of a canonical form and showing its relation with  $\mathcal{N}_1$ -functions. Let  $H \in \mathbb{C}^{(n+1) \times (n+1)}$  ( $n \in \mathbb{N} \setminus \{0\}$ ) be an invertible, Hermitian-symmetric matrix. We say that  $X \in \mathbb{C}^{(n+1) \times (n+1)}$  is  *$H$ -selfadjoint* if  $X^* H = H X$ . Our main interest will lie in the matrix

$$(6) \quad H = \begin{bmatrix} -1 & 0 \\ 0 & I_n \end{bmatrix},$$

---

<sup>1</sup>For arbitrary  $\mathcal{N}_1$  function  $z_0 = \infty$  can be also the GZNT, in that case  $\lim_{z \rightarrow \infty} zQ(z) \in [0, \infty)$ . However, this is clearly not possible for  $Q$  of the form (3).

where  $I_n$  denotes the identity matrix of size  $n \times n$ . As it was already mentioned, each  $H$ -selfadjoint matrix has the form

$$(7) \quad X = \begin{bmatrix} a & -b^* \\ b & C \end{bmatrix},$$

with  $a \in \mathbb{R}$ ,  $b \in \mathbb{C}^n$  and hermitian-symmetric  $C \in \mathbb{C}^{n \times n}$ . Due to [14] there exists an invertible matrix  $S$  and a pair of matrices  $H', S' \in \mathbb{C}^{(n+1) \times (n+1)}$  such that  $X = S^{-1}X'S$ ,  $H = S'^*H'S$  and  $X', H'$  are of one of the following forms:

Case 1.

$$X' = \begin{bmatrix} \beta & 0 \\ 0 & \bar{\beta} \end{bmatrix} \oplus \text{diag}(\zeta_2, \dots, \zeta_n), \quad H' = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \oplus I_{n-1},$$

with  $\beta \in \mathbb{C}^+$ ,  $\zeta_2, \dots, \zeta_n \in \mathbb{R}$ .

Case 2.

$$X' = [\beta] \oplus \text{diag}(\zeta_1, \dots, \zeta_n), \quad H' = [-1] \oplus I_n,$$

with  $\beta \in \mathbb{R}$ ,  $\zeta_1, \dots, \zeta_n \in \mathbb{R}$ .

Case 3.

$$X' = \begin{bmatrix} \beta & 1 \\ 0 & \beta \end{bmatrix} \oplus \text{diag}(\zeta_2, \dots, \zeta_n), \quad H' = \gamma \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \oplus I_{n-1},$$

with  $\beta \in \mathbb{R}$ ,  $\zeta_2, \dots, \zeta_n \in \mathbb{R}$ ,  $\gamma \in \{-1, 1\}$ .

Case 4.

$$X' = \begin{bmatrix} \beta & 1 & 0 \\ 0 & \beta & 1 \\ 0 & 0 & \beta \end{bmatrix} \oplus \text{diag}(\zeta_3, \dots, \zeta_n), \quad H' = \begin{bmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{bmatrix} \oplus I_{n-2},$$

with  $\beta \in \mathbb{R}$ ,  $\zeta_3, \dots, \zeta_n \in \mathbb{R}$ .

It is easy to verify that in each case  $X'$  is  $H'$ -symmetric. The pair  $(X', H')$  is called *the canonical form* of  $(X, H)$ . We refer the reader to [14] for the proof and for canonical forms for general  $H$ -symmetric matrices and to [5, 20] for the infinite-dimensional counterpart of the theory. At this point is enough to mention that the canonical form is uniquely determined (up to permutations of the numbers  $\zeta_i$ ) for each pair  $(X, H)$ , where  $X$  is  $H$ -selfadjoint. Note that in each of the cases  $\beta$  is an eigenvalue of  $X$  and there exists a corresponding eigenvector  $x \in \mathbb{C}^{n+1}$  satisfying  $[x, x]_H \leq 0$ , furthermore,  $\beta$  is the only eigenvalue in  $\mathbb{C}^+ \cup \mathbb{R}$  having this property. Therefore, we will call  $\beta$  the *eigenvalue of nonpositive type* of  $X$ .

Observe that the function

$$(8) \quad Q(z) = a - z + b^*(C - z)^{-1}b$$

is an  $\mathcal{N}_1$ -function. Indeed, if  $D = UCU^* = \text{diag}(\lambda_1, \dots, \lambda_n)$  is a diagonalization of the hermitian-symmetric matrix  $C$  and  $d = Ub$  then

$$Q(z) = a - z + \sum_{j=1}^n \frac{|d_j|^2}{\lambda_j - z} = a - z + \hat{\mu}(z), \quad \text{where} \quad \mu = \sum_{j=1}^n |d_j|^2 \delta_{\lambda_j},$$

and we may apply Proposition 1. The following lemma is a standard in the indefinite linear algebra theory. We present the proof for the reader's convenience.

**Lemma 3.** *Let  $X$  and  $Q$  be defined by (7) and (8), respectively. A point  $\beta \in \mathbb{C}^+ \cup \mathbb{R}$  is the eigenvalue of nonpositive type of  $X$  if and only if it is the GZNT of  $Q(z)$ . Furthermore, the algebraic multiplicity of  $\beta$  as an eigenvalue of  $X$  equals the order of  $\beta$  as a zero of  $Q(z)$ .*

*Proof.* First note that, due to the Shur complement formula<sup>2</sup>,

$$-\frac{1}{Q(z)} = e^* H(X - z)^{-1} e,$$

where  $e$  denotes the first vector of the canonical basis of  $\mathbb{C}^{n+1}$ . Let  $(X', H')$  be the canonical form of  $(X, H)$  and let  $S$  be the appropriate transformation. Consequently,

$$(9) \quad -\frac{1}{Q(z)} = (Se_1)^* H'(X' - z)^{-1} Se_1.$$

Below we evaluate this expression in each of the Cases 1–4. Let  $f = [f_0, \dots, f_n]^\top = Se$ . Note that

$$(10) \quad f_1^* H' f_1 = e_1^* H e_1 = -1,$$

independently on the Case.

Case 1. Observe that  $f_0 \bar{f}_1 \neq 0$ , otherwise  $f^* H' f \geq 0$ , which contradicts (10). Due to (9) one has

$$-\frac{1}{Q(z)} = \frac{f_0 \bar{f}_1}{\beta - z} + \frac{f_1 \bar{f}_0}{\bar{\beta} - z} + \sum_{j=2}^n \frac{|f_j|^2}{\zeta_j - z}.$$

Hence,  $\beta \in \mathbb{C}^+$  is a simple pole of  $-1/Q$  and consequently it is the GZNT of  $Q$  and a simple zero of  $Q$ .

Case 2. Observe that  $|f_0|^2 > \sum_{j=1}^n |f_j|^2$ , otherwise  $f^* H' f \geq 0$ , which contradicts (10). Due to (9) one has

$$-\frac{1}{Q(z)} = \frac{-|f_0|^2}{\beta - z} + \sum_{j=1}^n \frac{|f_j|^2}{\zeta_j - z}.$$

Hence, the residue of  $-1/Q$  in  $\beta$  is less than zero. Consequently  $Q(\beta) = 0$ ,  $Q'(\beta) < 0$  and  $\beta$  is the GZNT of  $Q$ .

Case 3. Observe that  $|f_1|^2 > 0$ , otherwise  $f^* H' f \geq 0$ , which contradicts (10). Due to (9) one has

$$-\frac{1}{Q(z)} = \frac{2\gamma \operatorname{Re} f_0 \bar{f}_1}{\beta - z} + \frac{-\gamma |f_1|^2}{(\beta - z)^2} + \sum_{j=2}^n \frac{|f_j|^2}{\zeta_j - z}.$$

Hence,  $\beta$  is pole of  $-1/Q$  of order 2. Consequently,  $Q(\beta) = Q'(\beta) = 0$ ,  $Q''(\beta) \neq 0$  and  $\beta$  is the GZNT.

Case 4. Observe that  $|f_2|^2 > 0$ , otherwise  $f^* H' f \geq 0$ , which contradicts (10). Due to (9) one has

$$-\frac{1}{Q(z)} = \frac{2 \operatorname{Re} f_0 \bar{f}_2 + |f_1|^2}{\beta - z} + \frac{-2 \operatorname{Re} f_1 \bar{f}_2}{(\beta - z)^2} + \frac{|f_2|^2}{(\beta - z)^3} + \sum_{j=3}^n \frac{|f_j|^2}{\zeta_j - z},$$

Hence,  $\beta$  is pole of  $-1/Q$  of order 3. Consequently,  $Q(\beta) = Q'(\beta) = Q''(\beta) = 0$ ,  $Q'''(\beta) \neq 0$  and  $\beta$  is the GZNT of  $Q$ .  $\square$

---

<sup>2</sup>It is well known [17] that  $-1/Q$  belongs to  $\mathcal{N}_1$  provided that  $Q$  belongs to  $\mathcal{N}_1$ , however, this information is not essential for the proof.

3. RANDOM  $H$ -SELFADJOINT MATRICES

By  $X_N$ ,  $H_N$  we understand the following pair of a random and deterministic matrix in  $\mathbb{C}^{(N+1) \times (N+1)}$

$$(11) \quad X_N = \begin{bmatrix} a_N & -b_N^* \\ b_N & C_N \end{bmatrix}, \quad H_N = \begin{bmatrix} -1 & 0 \\ 0 & I_N \end{bmatrix},$$

where  $a_N$  is a real-valued random variable,  $b_N$  is a random vector in  $\mathbb{C}^N$ , and  $C_N$  is a hermitian-symmetric random matrix in  $\mathbb{C}^{N \times N}$ . Note that  $X_N$  is  $H_N$ -symmetric. By  $\lambda_1^N \leq \dots \leq \lambda_N^N$  we denote the eigenvalues of  $C_N$  and by  $\nu_N$  we denote the random measure on  $\mathbb{R}$

$$\nu_N = \frac{1}{N} \sum_{j=1}^N \delta_{\lambda_j^N}.$$

Recall that

$$(12) \quad \hat{\nu}_N(z) = \frac{\text{tr}(C_N - z)^{-1}}{N}.$$

The assumptions on  $X_N$  are as follows:

- (R0) The random variable  $a_N$  is independent on the entries of the vector  $b_N$  and on the entries of the matrix  $C_N$  for each  $N > 0$ , furthermore  $a_N$  converges with  $N \rightarrow \infty$  to zero in probability.
- (R1) The random vector  $b_N$  is of the form

$$b_N := \frac{1}{\sqrt{N}} [x_{j0}]_{j=1, \dots, N},$$

where  $[x_{j0}]_{j>0}$  are i.i.d. random variables, independent on the entries of  $C_N$  for  $N > 0$ , of zero mean with  $E|x_{j0}|^2 = s^2$  for  $j > 0$ .

- (R2) The random measure  $\nu_N$  converges with  $N \rightarrow \infty$  to some non-random measure  $\mu_0$  weakly in probability

All the results below hold also in the case when all variables  $x_{j0}$  ( $j > 0$ ) are real, in this situation  $b_N^*$  is just the transpose of  $b_N$ . The entries of  $C_N$  might be as well real or complex. In Section 5 we will consider two instances of the matrix  $C_N$ : a Wigner matrix and a diagonal matrix. In the case when  $C_N$  is a Wigner matrix the proposition below is a consequence of the isotropic semicircle law [12, 15]. We present below a simple proof of the general case, based on the ideas in [22].

**Proposition 4.** *Assume that (R1) and (R2) are satisfied. Then for each  $z \in \mathbb{C}^+$*

$$b_N^*(C_N - z)^{-1}b_N \rightarrow s^2 \hat{\mu}_0(z) \quad (N \rightarrow \infty)$$

*in probability.*

By  $\|y\|$  we denote the euclidean norm of  $y \in \mathbb{C}^n$ .

*Proof.* In the light of Chebyshev's inequality, (12) and assumption (R2) it is enough to show that

$$(13) \quad \lim_{N \rightarrow \infty} E \left| b_N^*(C_N - z)^{-1}b_N - s^2 \frac{\text{tr}(C_N - z)^{-1}}{N} \right|^2 = 0.$$

Observe that

$$(14) \quad E \left| b_N^*(C_N - z)^{-1}b_N - s^2 \frac{\text{tr}(C_N - z)^{-1}}{N} \right|^2 =$$

$$(15) \quad \left( E |b_N^*(C_N - z)^{-1} b_N|^2 - s^4 E \left| \frac{\text{tr}(C_N - z)^{-1}}{N} \right|^2 \right) - 2 \operatorname{Re} E \left( s^2 \frac{\overline{\text{tr}(C_N - z)^{-1}}}{N} \left( b_N^*(C_N - z)^{-1} b_N - s^2 \frac{\text{tr}(C_N - z)^{-1}}{N} \right) \right).$$

First we prove that the summand (15) equals zero. Indeed, conditioning on the  $\sigma$ -algebra generated by the entries of the matrix  $C_N$  and setting

$$[c_{ij}]_{i,j=1}^N = (C_N - z)^{-1}$$

one obtains

$$\begin{aligned} E \left( s^2 \frac{\overline{\text{tr}(C_N - z)^{-1}}}{N} \left( b_N^*(C_N - z)^{-1} b_N - s^2 \frac{\text{tr}(C_N - z)^{-1}}{N} \right) \right) = \\ E \left( s^2 \sum_{i=1}^N \frac{\overline{c_{ii}}}{N} \left( \sum_{j,k=1}^N c_{jk} \frac{x_{0j} \overline{x_{0k}}}{N} - s^2 \sum_{j=1}^N \frac{c_{jj}}{N} \right) \right) = \\ E \left( s^2 \sum_{i=1}^N \frac{\overline{c_{ii}}}{N} \left( \sum_{j=1}^N c_{jj} \frac{s^2}{N} - s^2 \sum_{j=1}^N \frac{c_{jj}}{N} \right) \right) = 0. \end{aligned}$$

Next, observe that

$$\begin{aligned} E |b_N^*(C_N - z)^{-1} b_N|^2 &= E \sum_{i,j,k,l=1}^N c_{ij} \overline{c_{kl}} \frac{x_{0i} \overline{x_{0j}} x_{0k} \overline{x_{0l}}}{N^2} = \\ s^4 \sum_{i,j=1}^N \frac{E(c_{ii} \overline{c_{jj}})}{N^2} + s^4 \sum_{i,j=1}^N \frac{E(c_{ij} \overline{c_{ij}})}{N^2} &= s^4 E \left| \frac{\text{tr}(C_N - z)^{-1}}{N} \right|^2 + s^4 E \sum_{i,j=1}^N \frac{c_{ij} \overline{c_{ij}}}{N^2}. \end{aligned}$$

This allows us to estimate (14) by

$$s^4 \left| E \sum_{i,j=1}^N \frac{c_{ij} \overline{c_{ij}}}{N^2} \right| \leq s^4 E \frac{\|(C_N - z)^{-1}\|^2}{N} = \frac{s^4 \operatorname{dist}(z, \sigma(C_N))^{-2}}{N} \leq \frac{s^4}{\operatorname{Im} z^2 N},$$

which finishes the proof of (13).  $\square$

Let  $U_N$  be a unitary matrix, such that  $U_N C_N U_N^*$  is diagonal and let  $d_N = [d_1^N, \dots, d_N^N]^\top = U_N b_N$ . Denote by  $\mu_N$  the measure defined by

$$\mu_N = \sum_{j=1}^N |d_j^N|^2 \delta_{\lambda_j^N},$$

and observe that  $\hat{\mu}_N(z) = b_N^*(C_N - z)^{-1} b_N$ .

**Proposition 5.** *Assume that (R1) and (R2) are satisfied. Then the sequence of random measures  $\mu_N$  converge weakly with  $N \rightarrow \infty$  to  $\mu_0$  in probability.*

*Proof.* First note that almost surely  $\mu_N(\mathbb{R}) \rightarrow s^2 \mu_0(\mathbb{R})$  with  $N \rightarrow \infty$ . Indeed,

$$\mu_N(\mathbb{R}) = \sum_{j=1}^N |d_j^N|^2 = \|d_N\|^2 = \|b_N\|^2 = \frac{1}{N} \sum_{j=1}^N |x_{0j}|^2,$$

which converges almost surely to  $s^2$  by the strong law of large numbers. Furthermore, Proposition 4 shows that  $\hat{\mu}_N(z)$  converges in probability to  $\hat{\mu}_0(z)$  for every  $z \in \mathbb{C}^+$ . Repeating the proof of Theorem 2.4.4 of [1] we get the weak convergence of  $\mu_N$  in probability.  $\square$

#### 4. MAIN RESULTS

**Theorem 6.** *If (R0) – (R2) are satisfied then the eigenvalue of nonpositive type  $\beta_N$  of  $X_N$  converges in probability to the GZNT  $\beta_0$  of the  $\mathcal{N}_1$ -function*

$$Q_0(z) = -z + s^2 \hat{\mu}_0(z).$$

*Proof.* Consider a sequence of  $\mathcal{N}_1$ -functions

$$(16) \quad Q_N(z) = a_N - z + \hat{\mu}_N(z).$$

Recall that each of those functions has precisely one GZNT which, by definition of  $\mu_N$  and Lemma 3, is the eigenvalue of nonpositive type  $\beta_N$  of  $X_N$ . Recall that  $a_N$  converges to zero in probability by (R0) and  $\mu_N$  converges to  $\mu_0$  in probability by Proposition 5. Let  $d$  be any metric that metrizes the topology of weak convergence on  $M_b^+(\mathbb{R})$ . Since  $\beta_N$  is a continuous function of  $\mu_N$  and  $a_N$  (Proposition 2), for each  $\varepsilon > 0$  one can find  $\delta > 0$  such that for each  $N > 0$  the event  $\{|a_N| < \delta, d(\mu_N, \mu_0) < \delta\}$  is contained in  $\{|\beta_N - \beta_0| < \varepsilon\}$ . Using the assumed in (R0) independence of  $\mu_N$  and  $a_N$  one obtains

$$P(|\beta_0 - \beta_N| \geq \varepsilon) \leq P(|a_N| \geq \delta) \cdot P(d(\mu_N, \mu_0) \geq \delta).$$

Hence,  $\beta_N$  converges to  $\beta_0$  in probability.  $\square$

As it was explained in Section 1, each matrix  $X_N$  has, besides the eigenvalue  $\beta_N$  of nonpositive type, a set of real eigenvalues  $\zeta_{k_N}^N, \dots, \zeta_N^N$ , where  $k_N = 1$  in Case 1 and 3,  $k_N = 2$  in Case 2 and  $k_N = 3$  in Case 3. By  $\tau_N$  we denote the empirical measure connected with these eigenvalues:

$$\tau_N = \frac{1}{N} \sum_{j=k_N}^N \delta_{\zeta_j^N}.$$

**Theorem 7.** *If (R0)–(R2) are satisfied and the random variables  $\{x_{0j} : j > 0\}$  are continuous, then the measure  $\tau_N$  converges weakly in probability to  $\mu_0$ .*

*Proof.* We use the notations  $U_N, d_N$  and  $\mu_N$  from the previous section, let also  $Q_N$  be given by (16). Note that the set  $\{y \in \mathbb{C}^N : (U_N y)_j = 0\}$  is of Lebesgue measure zero. Hence, with probability one  $d_j^N \neq 0$  for  $j = 1, \dots, N$ ,  $N > 0$ . Therefore,  $\hat{\mu}_N(z)$  is a rational function almost surely with poles of order one in  $\lambda_1^N, \dots, \lambda_N^N$ . Furthermore,

$$Q_N(z) = \frac{(a_N - z) \prod_{j=1}^N (\lambda_j^N - z) + \sum_{i=1}^N |d_i^N|^2 \prod_{j \neq i} (\lambda_j^N - z)}{\prod_{j=1}^N (\lambda_j^N - z)}.$$

In consequence,  $Q_N$  has exactly  $N + 1$  zeros counting multiplicities, all of them different from  $\lambda_1^N, \dots, \lambda_N^N$ . Due to the Schur complement argument, each of those

zeros is an eigenvalue of the matrix  $X_N \in \mathbb{C}^{(N+1) \times (N+1)}$ . Furthermore, due to Lemma 3 the algebraic multiplicity of  $\beta_N$  as eigenvalue of  $X_N$  equals the order of  $\beta_N$  as a zero of  $Q_N$ . In consequence, the spectrum of  $X_N$  coincides with the zeros of  $Q_N$  and  $\beta_N$  is the only zero of order possibly greater than one<sup>3</sup>.

On the other hand, the function  $\hat{\mu}_N$  is increasing on the real line with simple poles in  $\lambda_1^N, \dots, \lambda_N^N$ . Hence, in each of the intervals  $(\lambda_j^N, \lambda_{j+1}^N)$  ( $j = 1, \dots, N-1$ ) there is an odd number of zeros of  $Q_N$ , counting multiplicities. Consequently, in each of the intervals  $(\lambda_j^N, \lambda_{j+1}^N)$  ( $j = 1, \dots, N-1$ ) there is precisely one zero of  $Q_N$ , except possibly one interval that contains three zeros of  $Q_N$ . Out of these three zeros of  $Q_N$  either one or two of them belong to the set  $\{\zeta_{k_N}^N, \dots, \zeta_N^N\}$ , accordingly to the canonical form of  $X_N$ . Hence, in each of the intervals  $(\lambda_j^N, \lambda_{j+1}^N)$  ( $j = 1, \dots, N-1$ ) there is precisely one of the eigenvalues  $\zeta_{k_N}^N, \dots, \zeta_N^N$ , except possibly one interval that contains two of the eigenvalues  $\zeta_{k_N}^N, \dots, \zeta_N^N$ . Consequently, the weak limit of  $\tau_N$  in probability equals the weak limit of  $\nu_N$ .  $\square$

## 5. TWO INSTANCES

In the present section we consider two instances of  $C_N$ : the Wigner matrix and the diagonal matrix. These both cases appear naturally as applications of main results. We refer the reader to [25] for a scheme joining both examples.

Consider an  $H$ -selfadjoint real Wigner matrix

$$(17) \quad X_N := \frac{1}{\sqrt{N}} H_N [x_{ij}]_{i,j=0}^N,$$

with  $x_{ij}$  real,  $x_{ij} = x_{ji}$  ( $0 \leq i < j < \infty$ ), i.i.d., of zero mean and variance equal to  $s^2$ , and let  $H_N$  be defined as in (11). Clearly  $X_N$  is  $H_N$ -selfadjoint and satisfies (R0)–(R2) with  $\mu_0$  equal to the Wigner semicircle measure  $\sigma$ . The Stieltjes transform of the  $\sigma$  equals

$$\hat{\sigma}(z) = \frac{-z + \sqrt{z^2 - 4s^2}}{2s^2}.$$

It is easy to check that  $\beta_0 = \frac{\sqrt{2}}{2}s i$  is a zero of  $Q_0(z) = -z + s^2 \hat{\sigma}(z)$ . Hence,  $\beta_0$  is the GZNT of  $Q_0$  and we have proved the first part of the theorem below.

**Theorem 8.** *Let  $X_N$  be defined by (17). Then*

- (i)  $\beta_N$  converges in probability to  $\beta_0 = \frac{\sqrt{2}}{2}s i$ ;
- (ii) *if, additionally, the random variables  $b_{ij}$  ( $0 \leq i < j < \infty$ ) are continuous, then the probability of an event that there are precisely  $N-1$  real eigenvalues  $\zeta_2^N < \dots < \zeta_N^N$  of  $X_N$  and the inequalities*

$$(18) \quad \lambda_1^N < \zeta_2^N < \lambda_2^N < \dots < \lambda_{N-1}^N < \zeta_N^N < \lambda_N^N.$$

*are satisfied, converges with  $N$  to 1.*

*Proof.* (ii) Assume that

$$(19) \quad |\beta_N - \beta_0| \leq \frac{\sqrt{2}}{4}s.$$

Then the canonical form of  $(X_N, H_N)$  is as in Case 1. In consequence, there are exactly  $N-1$  real eigenvalues  $\zeta_2^N, \dots, \zeta_N^N$  of  $X_N$ . Let us recall now the arguments

---

<sup>3</sup>In other words:  $e$  is almost surely a cyclic vector of  $X_N$ .





FIGURE 2. The real and imaginary part of  $\beta_N$ , with real, gaussian entries of  $X_N$  and  $s^2 = 1$ , computed with R [28].

from proof of Theorem 7. The function  $\hat{\mu}_N$  is increasing on the real line with simple poles in  $\lambda_1^N, \dots, \lambda_N^N$ . In each of the intervals  $(\lambda_j^N, \lambda_{j+1}^N)$  ( $j = 1, \dots, N-1$ ) there is at least one of the eigenvalues  $\zeta_2^N, \dots, \zeta_N^N$ . Consequently, each of the intervals  $(\lambda_j^N, \lambda_{j+1}^N)$  ( $j = 1, \dots, N-1$ ) contains precisely one of the eigenvalues  $\zeta_2^N, \dots, \zeta_N^N$ . To finish the proof it is enough to note that by point (i) for every  $\varepsilon > 0$  there exists  $N_0 > 0$  such that for  $N > N_0$  the probability of (19) is greater than  $1 - \varepsilon$ .  $\square$

The numerical simulations of values of  $\text{Re } \beta_N$  and  $\text{Im } \beta_N$  can be found in Figure 2. Note that  $\beta_0$  lies in open upper half-plane and (4) is satisfied. We provide now an example when  $\beta_0 \in \mathbb{R}$  and show that each number in  $[0, \infty)$  can be the limit in (5). Let  $a_N = 0$ ,  $x_{i0}$  ( $i = 1, 2, \dots$ ) be independent real variables of zero mean and variance  $s^2$  and let  $C_N = \text{diag}(c_1, \dots, c_N)$ , where the random variables  $\{c_j : j = 1, \dots\}$  are i.i.d. and independent on  $x_{i0}$  ( $i = 1, 2, \dots$ ). Furthermore, let

FIGURE 3. The imaginary part of  $\beta_N$ .

the law of  $c_j$  (which is simultaneously the limit measure  $\mu_0$ ) be given by a density

$$\phi(t) = \begin{cases} \frac{3t^2}{2} & : t \in [-1, 1] \\ 0 & : t \in \mathbb{R} \setminus [-1, 1] \end{cases}.$$

An easy calculation shows that

$$\lim_{z \rightarrow 0} \frac{\hat{\mu}_0(z)}{z} = 3.$$

Hence,

$$\lim_{z \rightarrow 0} \frac{-z + s^2 \hat{\mu}_0(z)}{z} = -1 + 3s^2$$

and the function

$$Q_0(z) = -z + \hat{\mu}_0(z) = -z + \int_{\mathbb{R}} \frac{\phi(t)}{t - z} dt$$

has a GZNT at  $z = 0$  if  $s^2 \leq 1/3$ . Note that  $\beta_0 = 0$  lies in the support of  $\mu_0$ . The case  $s^2 = 1/3$  is plotted in Figure 3. Only the imaginary part is displayed, since the numerical computation of the real part of  $\beta_N$  might be not reliable in case  $\beta_N \in \mathbb{R}$ . One may observe that the convergence of  $\beta_N$  is worse in Figure 2. Also, the canonical form of  $(X_N, H_N)$  changes with  $N$ , contrary to the case when  $H_N X_N$  is a Wigner matrix. In the case  $s^2 < 1/3$  in numerical simulations point  $\beta_N$  is real for all  $N$ .

#### REFERENCES

- [1] G.W. Anderson, A. Guionnet, O. Zeitouni, *An Introduction to Random Matrices*, Cambridge University Press 2010.
- [2] F. Benaych-Georges, A. Guionnet, and M. Maïda, Fluctuations of the extreme eigenvalues of finite rank deformations of random matrices, arXiv:1009.0145.
- [3] F. Benaych-Georges, A. Guionnet, and M. Maïda, Large deviations of the extreme eigenvalues of random deformations of matrices, *Prob. Theor. Rel. Fields* (2010), 1–49.

- [4] F. Benaych-Georges and R.R. Nadakuditi, The eigenvalues and eigenvectors of finite, low rank perturbations of large random matrices, *Adv. Math.* 227 (2011), 494–521.
- [5] J. Bognár, *Indefinite Inner Product Spaces*, Springer-Verlag, New York-Heidelberg, 1974.
- [6] M. Capitaine, C. Donati-Martin, and D. Féral, The largest eigenvalues of finite rank deformation of large Wigner matrices: convergence and nonuniversality of the fluctuations, *Ann. Prob.* 37 (2009), 1–47.
- [7] M. Capitaine, C. Donati-Martin, D. Féral and M. Février, Free convolution with a semi-circular distribution and eigenvalues of spiked deformations of Wigner matrices, *Electronic Journal of Probability*, 16 (2011), 1750–1792.
- [8] V. Derkach, S. Hassi, and H.S.V. de Snoo, Operator models associated with Kac subclasses of generalized Nevanlinna functions, *Methods of Functional Analysis and Topology*, 5 (1999), 65–87.
- [9] V.A. Derkach, S. Hassi, and H.S.V. de Snoo, Rank one perturbations in a Pontryagin space with one negative square, *J. Functional Analysis*, 188 (2002), 317–349.
- [10] A. Dijksma, H. Langer, A. Luger, and Yu. Shondin, A factorization result for generalized Nevanlinna functions of the class  $\mathbf{N}_\kappa$ , *Integral Equations Operator Theory*, 36 (2000), 121–125.
- [11] W.F. Donoghue, *Monotone matrix functions and analytic continuation*, Springer-Verlag, New York-Heidelberg, 1974.
- [12] L. Erdős, Rigidity of eigenvalues of generalized Wigner matrices, *Adv. Math. Preprint* 229 (2012), 1435–1515.
- [13] D. Féral and S. Péché, The largest eigenvalue of rank one deformation of large Wigner matrices, *Comm. Math. Phys.* 272 (2007), 185–228.
- [14] I. Gohberg, P. Lancaster and L. Rodman: *Indefinite Linear Algebra and Applications*. Birkhäuser-Verlag, 2005.
- [15] A. Knowles, J. Yin, The Isotropic Semicircle Law and Deformation of Wigner Matrices, arXiv:1110.6449v2.
- [16] M.G. Kreĭn and H. Langer, The defect subspaces and generalized resolvents of a Hermitian operator in the space  $\Pi_\kappa$ , *Funkcional. Anal. i Priložen.* 5 (1971), 54–69 [Russian].
- [17] M.G. Kreĭn and H. Langer, Über einige Fortsetzungsprobleme, die eng mit der Theorie hermitescher operatoren im Raume  $\Pi_\kappa$  zusammenhangen. I. Einige Funktionenklassen und ihre Darstellungen, *Math. Nachr.*, 77 (1977), 187–236.
- [18] M.G. Kreĭn and H. Langer, Some propositions on analytic matrix functions related to the theory of operators in the space  $\Pi_\kappa$ , *Acta Sci. Math.* (Szeged), 43 (1981), 181–205.
- [19] H. Langer, A characterization of generalized zeros of negative type of functions of the class  $\mathbf{N}_\kappa$ , *Oper. Theory Adv. Appl.*, 17 (1986), 201–212.
- [20] I.S. Iohvidov, M.G. Krein, H. Langer, *Introduction to spectral theory of operators in spaces with indefinite metric*, Mathematical Research, vol. 9. Akademie-Verlag, Berlin, 1982.
- [21] H. Langer, A. Luger, V. Matsaev, Convergence of generalized Nevanlinna functions, *Acta Sci. Math.* (Szeged), 77 (2011), 425–437.
- [22] M. A. Marchenko, L. A. Pastur, Distribution of eigenvalues in some ensembles of random matrices, *Mat. Sb.*, 72(114) (1967), 507–536 [Russian].
- [23] S. N. Chandler-Wilde, R. Chonchaiya, M. Lindner, Eigenvalue problem meets Sierpinski triangle: computing the spectrum of a non-self-adjoint random operator, *Oper. Matrices* 5 (2011), 633648.
- [24] S. N. Chandler-Wilde, R. Chonchaiya, M. Lindner, On the Spectra and Pseudospectra of a Class of Non-Self-Adjoint Random Matrices and Operators, arXiv:1107.0177.
- [25] L. A. Pastur, On the spectrum of random matrices, *Theoretical and Mathematical Physics*, 10 (1972), 102–112, [Russian].
- [26] A. Pizzo, D. Renfrew, and A. Soshnikov, On finite rank deformations of Wigner matrices, to appear in *Ann. Inst. Henri Poincaré (B)*, arXiv:1103.3731.
- [27] L.S. Pontryagin, Hermitian operators in spaces with indefinite metric, *Izv. Nauk. Akad. SSSR, Ser. Math.* 8 (1944), 243–280 [Russian].
- [28] R Development Core Team (2011). R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. ISBN 3-900051-07-0, URL <http://www.R-project.org/>.
- [29] H.S.V. de Snoo, H. Winkler, M. Wojtylak, Zeros and poles of nonpositive type of Nevanlinna functions with one negative square, *J. Math. Anal. Appl.*, 382 (2011), 399–417.

- [30] R. Vershynin, Spectral norm of products of random and deterministic matrices, *Probab. Theory Related Fields* 150 (2011), 471–509.
- [31] E. P. Wigner. Characteristic vectors of bordered matrices with infinite dimensions. *Annals Math.*, 62(1955), 548–564.

FACULTY OF MATHEMATICS AND COMPUTER SCIENCE, JAGIELLONIAN UNIVERSITY, ŁOJASIEWICZA  
6, 30-348 KRAKÓW, POLAND  
*E-mail address:* `michal.wojtylak@gmail.com`